Understanding the relationship between resonant and non-resonant x-ray probes of correlated matter

C. J. Jia,[1, 2] C.-C. Chen,[1, 3] A. P. Sorini,[1] B. Moritz,[1, 4] and T. P. Devereaux[1, 5]

[1]*Stanford Institute for Materials and Energy Sciences,*
SLAC National Accelerator Laboratory, 2575 Sand Hill Road, Menlo Park, CA 94025.
[2]*Departments of Applied Physics, Stanford University, Stanford, CA 94305.*
[3]*Departments of Physics, Stanford University, Stanford, CA 94305.*
[4]*Department of Physics and Astrophysics, University of North Dakota, Grand Forks, ND 58202, USA*
[5]*Geballe Laboratory for Advanced Materials, Stanford University, Stanford, CA 94305.*

(Dated: September 16, 2011)

arXiv:1109.3446v1 [cond-mat.str-el] 15 Sep 2011

A great deal of effort has been devoted to understanding the possible connections between resonant and non-resonant x-ray probes of correlated materials. In this letter we present a comparison between momentum-dependent indirect resonant inelastic x-ray scattering (RIXS) and the dynamical structure factor $S(\mathbf{q},\omega)$ for the single-band Hubbard model. Our results highlight the fact that RIXS exhibits charge as well as spin or multi-magnon energy-loss excitations with a strong incident energy dependence. Although indirect RIXS and $S(\mathbf{q},\omega)$ share a number of the same possible final state excitations, the cross-sections as functions of energy-loss are quite different; and the two spectra can not always be simply connected by resonant prefactors or by other commonly proposed approximation schemes.

PACS numbers: 78.70.Ck, 74.72 -h, 78.20 Bh

Resonant x-ray probes, such as resonant elastic and inelastic x-ray scattering (REXS and RIXS)[1, 2], resonantly excite core electrons when tuning the incident photon energy to a particular absorption edge. The resulting intermediate core-hole state could connect many-body ground state and final excited states via pathways that may be inaccessible via other non-resonant x-ray probes, such as nonresonant x-ray and electron scattering which are related to the charge dynamical structure factor $S(\mathbf{q},\omega)$. Tuning the incident photon energy to highlight different intermediate state can resonantly enhance certain subdominant characters of ground state wavefunction and final state excitations.[3] In this regard, resonance probe provides more information about dynamics and excitations than the non-resonant response.

Especially in correlated materials, resonance probes can provide wealth of information about charge, spin and orbital degree of freedom beyond the capability of non-resonant probes. However, this also has made RIXS difficult to interpret, because the intermediate core-hole state increases the complexity of the problem from an effective two-particle process as in the non-resonant response to an effective four-particle process. Researchers have proposed that under certain approximations such as small[4] and large core-hole potentials[5], RIXS could be connected to $S(\mathbf{q},\omega)$ by some resonant prefactor[4–6]. Experimentally, it has been shown that RIXS, or the extracted response function from RIXS, shares similar structure to $S(\mathbf{q},\omega)$ [7–9]. However there is no definitive proof that this connection works when the intermediate state core-hole potential is comparable to other energy scales in the system, especially at transition metal K-edges, like that for indirect RIXS at the Cu K-edge in cuprates where the core-hole potential can be on the same order as the Coulomb interaction and other energy scales.

In this work, we compare RIXS and $S(\mathbf{q},\omega)$ utilizing exact diagonalization and bi-conjugate gradient stabilized methods[10], to treat both response functions on equal footing within the intermediate core-hole potential regime without further approximation. Our study yields several important results. First, RIXS generally cannot be connected simply to $S(\mathbf{q},\omega)$, by a resonant prefactor within the intermediate core-hole potential regime. Second, for particular incident energies, corresponding to specific intermediate state configurations without appreciable core-hole potential "shake-up", RIXS and $S(\mathbf{q},\omega)$ can be connected well by a simple resonant prefactor. Third, we find two-magnon excitations for all the momentum points accessible in our calculations except at (π,π). The two-magnon peak at $(0,0)$ is consistent with theoretical prediction[11], but has not been observed experimentally, however these results may help to resolve this controversy given recent improvements in experimental resolution[2].

The single-band Hubbard Hamiltonian that serves as an effective low energy model for the cuprates can be written as

$$H = -\sum_{\mathbf{i},\mathbf{j}} t_{\mathbf{i},\mathbf{j}} d^{\dagger}_{\mathbf{i},\sigma} d_{\mathbf{j},\sigma} + \sum_{\mathbf{i}} U n^{d}_{\mathbf{i},\uparrow} n^{d}_{\mathbf{i},\downarrow}, \qquad (1)$$

where $d^{\dagger}_{\mathbf{i},\sigma}$ ($d_{\mathbf{i},\sigma}$) creates (annihilates) a fermion with spin σ on lattice site $\mathbf{i}$, $t_{\mathbf{i},\mathbf{j}}$ is the hopping integral restricted to the nearest- and next-nearest-neighbor, denoted as t and t'. U is the on-site Coulomb repulsion, and $n^{d}_{\mathbf{i},\sigma}$ is the fermion number operator. The momentum-dependent indirect RIXS cross section can be expressed using the

Kramers-Heisenberg formula as

$$I(\mathbf{q},\Omega,\omega_i) = \frac{1}{\pi}\mathrm{Im}\langle\Psi|\frac{1}{H-E_0-\Omega-i0^+}|\Psi\rangle \qquad (2)$$

where $|\Psi\rangle = \sum_{\mathbf{i},\sigma} e^{i\mathbf{q}\cdot\mathbf{r}_\mathbf{i}} D^\dagger \frac{1}{H'-E_0-\omega_i-i\Gamma} D|G\rangle$; $\mathbf{q}$ is the momentum transfer; ω_i and $\Omega = \omega_i - \omega_f$ are the incident energy and energy transfer, respectively; Γ is the inverse core-hole lifetime; E_0 is the ground state energy of the system absent the core-hole; $|G\rangle$ is the ground state wave function; $H' = H - U_c \sum'_{\mathbf{i},\sigma,\sigma} n^d_{\mathbf{i}\sigma} n^s_{\mathbf{i},\sigma'}$ represents the intermediate state Hamiltonian which includes the core-hole potential with strength U_c, with $n^s_{\mathbf{i},\sigma}$ denoting the core-hole number operator; D is the dipole transition operator, which for example in Cu K-edge RIXS excites a Cu $1s$ core electron to $4p$ level. The elastic lines will be removed for our RIXS calculations.

We also investigate the x-ray absorption spectroscopy (XAS) given by

$$B(\omega_i) = \frac{1}{\pi}\mathrm{Im}\langle G|D^\dagger\frac{1}{H'-E_0-\omega_i-i\Gamma}D|G\rangle \qquad (3)$$

and the charge dynamical structure factor given by

$$S(\mathbf{q},\omega) = \frac{1}{\pi}\mathrm{Im}\langle G|\rho^d_{-\mathbf{q}}\frac{1}{H-E_0-\omega-i0^+}\rho^d_{\mathbf{q}}|G\rangle, \qquad (4)$$

where $\rho^d_\mathbf{q} = \sum_{\mathbf{k},\sigma} d^\dagger_{\mathbf{k}+\mathbf{q},\sigma} d_{\mathbf{k},\sigma} = \sum_{\mathbf{i},\sigma} e^{i\mathbf{q}\cdot\mathbf{R}_\mathbf{i}} n^d_{\mathbf{i},\sigma}$.

We implement exact diagonalization calculations using parameters taken from the analysis of ARPES experiments in the cuprates.[12] The details of the calculations and parameter values are described in Ref. 13. Figure 1 shows the momentum-dependent indirect RIXS spectra for the half-filled, 12.5% hole-doped and 12.5% electron-doped single-band Hubbard model. At all dopings, RIXS displays strong resonances at the peaks of XAS (see the top row in Fig. 1). The resonance at incident energy $\sim -20t$ corresponds to an intermediate state where two electrons are bound at the core-hole site, which lowers the energy to $\sim -2U_c + U$. The resonance at incident energy $\sim -14t$ corresponds to a single electron bound at the core-hole site with energy $\sim -U_c$.[14, 15] These two states are typically denoted the well- and poorly-screened intermediate states, respectively. In both processes, the electrons experience a strong "shake-up" by the intermediate state core-hole potential via screening. On the other hand, the intermediate state corresponding to an incident energy $\sim 3t$ for the 12.5% hole-doped or $\sim -28t$ for the 12.5% electron-doped system, represents the state in which the core-hole has been created at an empty or a doubly-occupied site, respectively. In both cases the core-hole potential does not appreciably "shake-up" the electronic system.

We next examine the energy loss dependence in Fig. 1. For half-filling (Fig. 1(b)-(d)), RIXS displays a spectrum ranging from $\sim 5t$ up to $\sim 15t$ in energy loss, signaling excitations from the lower Hubbard band (LHB) to upper Hubbard band (UHB), which is mainly controlled by Coulomb U. The main peaks have strong momentum dependence with the shift to higher loss energy at larger momentum transfer. These features are consistent with various experiments[4, 7, 16, 17] and calculations[3, 14] carried out at the Cu K-edge in cuprates.



FIG. 1: (a)-(d) XAS and momentum-dependent indirect RIXS cross-section at momentum (0,0), (π,π) and $(0,\pi)$ at half-filling in the single-band Hubbard model. The blue arrows highlight two-magnon peaks in the RIXS cross-section. (f)-(h) the same for 12.5% hole-doping and (j)-(l) 12.5% electron-doping, respectively.

The RIXS spectra also exhibit distinctive structures at $\sim 1t$, highlighted by blue arrows in Fig. 1, emphasizing excitations of the spin degree of freedom - two-magnon excitations[11, 18–22]. Single-magnon excitations are forbidden in this case due to spin conservation, as we have neglected spin-orbit coupling. A strong two-magnon peak appears for momentum transfers of $(\pi, 0)$; however, no two-magnon low energy peaks appear at (π,π), as reported in recent K-edge RIXS measurements in LCO.[18, 19] However, we obtain two-magnon excitations at momentum transfer $(0, 0)$, which are symmetry allowed by the core-hole intermediate states. Similar



FIG. 2: Momentum-dependent RIXS and $S(\mathbf{q},\omega)$ in the half-filled single-band Hubbard model highlighting the (a) well-screened intermediate state at incident energy $E_{in} = -21.2t$ and (b) poorly-screened intermediate states at $E_{in} = -13.85t$. (c) RIXS at the well-screened intermediate state incident energy [from (a)] vs. $S(\mathbf{q},\omega)$ multiplied by a Lorentzian prefactor given by $((\omega_f - \omega_{res})^2 + \Gamma_f^2)^{-1}((\omega_i - \omega_{res} - |U|)^2 + \Gamma_i^2)^{-1}$, where ω_{res} is a free parameter.[7] We consider equal incident/outgoing photon lifetimes ($\Gamma_i = \Gamma_f$) for fitting. Note that the elastic lines of $S(\mathbf{q} = \mathbf{0},\omega)$ have been removed.

findings had also been reported in early calculations[14], however no experimental confirmation of the two-magnon feature at momentum $(0,0)$ has been reported, although recent improvements in resolution may resolve this issue.

When hole-doped and electron-doped (Figs. 1(f)-(h)and 1(j)-(l), respectively), the well-screened intermediate state mainly contains excitations across the Mott-gap, while the poorly-screened intermediate state show excitations both within the LHB and across the Mott-gap. This is consistent with recent intermediate state density of states calculations.[15] In the doped system low-energy charge and spin excitations should be located within the same energy-loss range, making it hard to distinguish between the two excitations. For the case in which the core-hole is created at an empty or a doubly-occupied site, the RIXS spectra only exhibit excitations within the LHB.

To make comparisons between RIXS and $S(\mathbf{q},\omega)$, in Fig. 2 we plot the RIXS spectra at intermediate states with the core-hole well- and poorly-screened, compared to $S(\mathbf{q},\omega)$ at half-filling. When $\omega > 5t$, $S(\mathbf{q},\omega)$ (black dotted) and the RIXS spectra (colored solid) generally span the same energy range at a given momentum transfer, and both shift to higher energy at larger momentum transfer. However, their overall intensities are very different: at momentum $(0,\pi/2)$, $(0,\pi)$ and $(\pi,\pi/2)$ the RIXS spectra have more weight at lower energy and a lower threshold; at momentum (π,π) and $(\pi/2,\pi/2)$ the RIXS spectra and $S(\mathbf{q},\omega)$ exhibit different main peaks and weights. Another striking difference lies in the $(0,0)$ momentum transfer. For $S(\mathbf{q},\omega)$ no energy loss excitations are allowed since the charge operator $\rho^d_{\mathbf{q}=\mathbf{0}}$ (see Eq. 3) commutes with the Hamiltonian; however, various excitations exist in RIXS and there is no such symmetry that forbids excitations. Moreover, RIXS spectra exhibit two-magnon excitations at low energy may be similar to the excitations revealed by the spin structure factor[6]. To investigate the conjecture that RIXS and $S(\mathbf{q},\omega)$ can be connected by a simple resonant prefactor, we compare RIXS at the well-screened intermediate state with $S(\mathbf{q},\omega)$ multiplied by a fitted Lorentzian prefactor as shown in Fig. 2(c). These results show that these two spectra only agree at momentum $(0,\pi)$ and $(\pi/2,\pi/2)$, thus a simple Lorentzian prefactor can not in general address the complicated differences between the spectra. The above analysis from half-filling also applies to other dopings for the poorly- and well-screened intermediate states, as shown in Fig. 3(a)-(b)and(d)-(e).

We find that in some limited cases RIXS and $S(\mathbf{q},\omega)$ agree well with one another, specifically for charge excitations within the LHB. This happens when the core-hole is created at an empty [Fig. 3(c)] or doubly-occupied site [Fig. 3(f)] for certain intermediate states, in which case the core-hole does not significantly "shake-up" the electrons, and the effect of the intermediate core-hole state can be neglected or is less prominent. Thus the complicated four-particle RIXS process could be simplified to the two-particle $S(\mathbf{q},\omega)$ process. Our results show that if the core-hole "shakes-up" the electrons when U_c is comparable to the other energy scales, RIXS deviates from $S(\mathbf{q},\omega)$; if there is a lack of this "shake-up" or core-hole screening, when U_c is extremely large or small when compared to the other energy scales, or for particular intermediate states, RIXS may be approximated by $S(\mathbf{q},\omega)$ with a resonant prefactor. In other words, the "shake-up" or screening is the key factor which differentiates RIXS from $S(\mathbf{q},\omega)$ in the range of intermediate value of U_c, consistent with other theoretical results.[15]

In summary, we have employed exact diagonalization to study indirect RIXS in the half-filled, hole- and electron-doped single-band Hubbard model over a wide range of incident energies and compared it to $S(\mathbf{q},\omega)$. In



FIG. 3: Momentum dependent Cu K-edge RIXS and $S(\mathbf{q},\omega)$ at 12.5% hole-doping [(a)$E_{in} = -21.2t$ (b)$E_{in} = -14.2t$ (c)$E_{in} = 3.78t$] and 12.5% electron-doping [(d)$E_{in} = -19.75t$ (e)$E_{in} = -13.33t$ (f)$E_{in} = -26.88t$]. We note that the elastic lines of $S(\mathbf{q} = \mathbf{0}, \omega)$ have been removed. Blue, red and purple lines represent RIXS spectra with the intermediate state having the core-hole well-screened, poorly-screened and created at an empty/doubly-occupied site, respectively.

general, the RIXS spectra show two-magnon excitations and exhibit inelastic excitations at momentum transfer $(0, 0)$, not captured by $S(\mathbf{q},\omega)$. The RIXS spectra at well-screened and poorly-screened intermediate states and $S(\mathbf{q},\omega)$ are significantly different for charge excitations both within the LHB and across the Mott-gap. In contrast, RIXS at empty or doubly-occupied intermediate states for various dopings reproduce $S(\mathbf{q},\omega)$ for excitations within the LHB. Our calculations demonstrate that although indirect RIXS has support on the same set of final state charge excitations as $S(\mathbf{q},\omega)$, the weights associated with each response function for each excitation are quite different and the two spectra generally could not be correlated by a simple prefactor. We further conclude that whether RIXS and $S(\mathbf{q},\omega)$ share similar weight depends strongly on the incident photon energy implying the importance of the intermediate state in RIXS; and the coincidence of RIXS and $S(\mathbf{q},\omega)$ is related to the lack of significant core-hole "shake-up" at these intermediate states.

This work was supported at SLAC and Stanford University by the U.S. Department of Energy, Office of Basic Energy Sciences, Division of Materials Science and Engineering, under Contract No. DE-AC02-76SF00515 and by the Computational Materials and Chemical Sciences Network (CMCSN) under Contract No. DE-FG02-08ER46540. C.J.Jia is also supported by the Stanford Graduate Fellows in Science and Engineering. A portion of the computational work was performed using the resources of the National Energy Research Scientific Computing Center (NERSC) supported by the U.S. Department of Energy, Office of Science, under Contract No. DE-AC02-05CH11231.

[1] A. Kotani and S. Shin, Rev. Mod. Phys. **73**, 203 (2001).
[2] L. J. P. Ament, M. van Veenendaal, T. P. Devereaux, J. P. Hill, and J. van den Brink, Rev. Mod. Phys **83**, 705 (2011).
[3] C.-C. Chen *et al*, Phys. Rev. Lett. **105**, 177401 (2010).
[4] P. Abbamonte *et al*, Phys. Rev. Lett. **83**, 860 (1999).
[5] J. van den Brink and M. v. Veenendaal, Europhys. Lett. **73**, 121 (2006).
[6] L. J. P. Ament *et al*, Phys. Rev. B **75**, 115118 (2007).
[7] J. Kim *et al*, Phys. Rev. B **79**, 094525 (2009).
[8] Y. J. Kim *et al*, Phys. Rev. Lett. **92**, 137402 (2004).
[9] Y. J. Kim *et al*, Phys. Rev. B **76**, 155116 (2007).
[10] H. A. Van der Vorst, SIAM Journal on Scientific and Statistical Computing **13**, 631 (1992).
[11] F. H. Vernay *et al*, Phys. Rev. B **75**, 020403(R) (2007).
[12] C. Kim *et al*, Phys. Rev. Lett. **80**, 4245 (1998).
[13] $U = 10t$, $t' = -0.34t$, $U_c = 15t$ and $\Gamma = t$, with $t = 0.35$eV. The calculations are done on the 16B Betts cluster (16-site square cluster) with periodic boundary conditions. The ground state wave function is obtained using the exact diagonalization (ED) technique utilizing Parallel Arnoldi PACKage (PARPACK), $|\Psi\rangle$ is calculated using the bi-conjugate gradient stabilized method (BiCGSTAB) and the final spectrum is calculated using continued fraction expansion method.
[14] K. Tsutsui *et al*, Phys. Rev. Lett. **83**, 3705 (1999).
[15] K. H. Anh *et al*, Phys. Rev. B **79**, 045103 (2009).
[16] G. Doring *et al*, Phys. Rev. B **70**, 085115 (2004).
[17] L. Lu *et al*, Phys. Rev. B **74**, 224509 (2006).
[18] J. P. Hill *et al*, Phys. Rev. Lett. **100**, 097001 (2008).
[19] D. S. Ellis *et al*, Phys. Rev. B **81**, 085124 (2010).
[20] F. Forte *et al*, Phys. Rev. B **77**, 134428 (2008).
[21] T. Nagao *et al*, Phys. Rev. B **75**, 214414 (2007).
[22] T. Tohyama, H. Onodera, K. Tsutsui, and S. Maekawa, Phys. Rev. Letts. **89**, 257405 (2002).